

**MEDICATION GUIDE**  
**ULORIC (Ū-'lor-ik)**  
**(febuxostat)**  
**tablets, for oral use**

Read the Medication Guide that comes with ULORIC before you start taking it and each time you get a refill. There may be new information. The Medication Guide does not take the place of talking with your doctor about your medical condition or your treatment.

**What is the most important information that I should know about ULORIC?**

**ULORIC may cause serious side effects, including:**

**Heart -related deaths.**

**Call your doctor or get emergency medical help right away if you have any of the following symptoms, especially if they are new, worse, or worry you:**

- chest pain
- shortness of breath or trouble breathing
- dizziness, fainting or feeling lightheaded
- rapid or irregular heartbeat
- numbness or weakness in one side of your body
- slurring of speech
- sudden blurry vision or sudden severe headache

**What is ULORIC?**

ULORIC is a prescription medicine called a xanthine oxidase (XO) inhibitor used to lower blood uric acid levels in adult patients with gout when allopurinol has not worked well enough or when allopurinol is not right for you. ULORIC is not for use in people who do not have symptoms of high blood uric acid levels. It is not known if ULORIC is safe and effective in children.

**Who should not take ULORIC?**

**Do not take ULORIC if you:**

- take azathioprine (Azasan, Imuran)
- take mercaptopurine (Purinethol, Purixan)

**What should I tell my doctor before taking ULORIC?**

**Before taking ULORIC tell your doctor about all of your medical conditions, including if you:**

- have taken allopurinol and what happened to you while you were taking it.
- have a history of heart disease or stroke.
- have liver or kidney problems.
- are pregnant or plan to become pregnant. It is not known if ULORIC will harm your unborn baby. Talk with your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if ULORIC passes into your breast milk. You and your doctor should decide if you should take ULORIC while breastfeeding.

**Tell your doctor about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. ULORIC may affect the way other medicines work, and other medicines may affect how ULORIC works.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

**How should I take ULORIC?**

- Take ULORIC exactly as your doctor tells you to take it.
- ULORIC can be taken with or without food.
- ULORIC can be taken with antacids.
- Your gout may get worse (flare) when you start taking ULORIC. **Do not stop taking ULORIC because you have a flare.**

Your doctor may do certain tests while you take ULORIC.

**What are the possible side effects of ULORIC?**

**ULORIC may cause serious side effects, including:**

- **Heart problems.** See “**What is the most important information I should know about ULORIC?**”.
- **Gout Flares.** Gout flares can happen when you start taking ULORIC. Your doctor may give you other medicines to help prevent your gout flares.
- **Liver problems.** Liver problems can happen in people who take ULORIC. Your doctor may do blood tests to check how well your liver is working before and during your treatment with ULORIC. Tell your doctor if you get any of the following signs or symptoms of liver problems:
  - fatigue
  - loss of appetite for several days or longer
  - pain, aching, or tenderness on the right side
  - dark or “tea-colored” urine
  - your skin or the white part of your eyes turns yellow (jaundice)

of your stomach-area

- **Severe skin and allergic reactions.** Serious skin and allergic reactions that may affect different parts of the body such as your liver, kidneys, heart or lungs, can happen in people who take ULORIC. Call your doctor right away or get emergency medical help if you have any of the following symptoms:
  - rash
  - red and painful skin
  - severe skin blisters
  - peeling skin
  - sores around the lips, eyes or mouth
  - swollen face, lips, mouth, tongue or throat
  - flu-like symptoms

**The most common side effects of ULORIC include:**

- abnormal liver function tests
- nausea
- joint pain
- rash

These are not all of the possible side effects of ULORIC.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**How should I store ULORIC?**

- Store ULORIC at room temperature.
- Keep ULORIC out of the light.

**Keep ULORIC and all medicines out of the reach of children.**

**General information about the safe and effective use of ULORIC.**

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use ULORIC for a condition for which it was not prescribed. Do not give ULORIC to other people, even if they have the same symptoms that you have. It may harm them.

You can ask your doctor or pharmacist for information about ULORIC that is written for health professionals.

**What are the ingredients in ULORIC?**

**Active ingredient:** febuxostat

**Inactive ingredients:** lactose monohydrate, microcrystalline cellulose, hydroxypropyl cellulose, sodium croscarmellose, silicon dioxide, magnesium stearate, and Opadry II, green

Distributed by:

**Takeda Pharmaceuticals America, Inc.**

Deerfield, IL 60015

ULORIC is a registered trademark of Teijin Limited registered in the U.S. Patent and Trademark Office and used under license by Takeda Pharmaceuticals America, Inc.

All other trademarks are the property of their respective owners.

©2009-2019 Takeda Pharmaceuticals America, Inc.

For more information, go to [www.ULORIC.com](http://www.ULORIC.com) or call 1-877-TAKEDA (1-877-825-3327).